

COCKSFOOT (DACTYLIS GLOMERATA)

MUSKETIER

Maturity group: medium (5)

Listing: since 2013

Advantages of MUSKETIER

- » Lowest susceptibility to rust in German cocksfoot assortment
- > Fine leaved variety with a high palatability and digestibility
- > Excellent contributor in meadow grass mixtures

Variety assessment of MUSKETIER

| Mass growth at the beginning | | medium |
|--------------------------------|-------------------|---------------|
| Tendency to winter kill | | medium |
| Tendency to lodging | - no assessment - | |
| Susceptibility to rust | | low |
| Sward density | | medium |
| Dry matter yield, total | | low to medium |
| Dry matter yield, 1st cut | | medium |
| Dry matter yield, further cuts | | medium |

(1 = very low, very early / 9 = very strong, very high, very late) Source: according to the specifying variety list 2022 issued by the Bundessortenamt

Variety descriptions above are provided to the best of our knowledge. Despite all efforts, SAATZUCHT STEINACH GmbH & Co KG cannot guarantee repeatability of described assessments in any case. The presented results should provide an appropriate knowledge base for decision support and may not be an indicator of results you may obtain in your individual location, as local growing conditions, including soil type and weather may vary. SAATZUCHT STEINACH GmbH & Co KG excludes all liability for damages which may arise from use of this variety description.

SAATZUCHT STEINACH GmbH & Co KG

Wittelsbacherstraße 15 · 94377 Steinach · GERMANY Telefon +49 (0) 9428 94190 · Telefax +49 (0) 9428 941930 · info@saatzucht.de · www.saatzucht.de